

Applications of somatic cell nuclear transfer in goats

Ton Yoisungnern* and Ashit Kumar Paul

School of Biotechnology, Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand.

*Corresponding author's e-mail: tonton9214@gmail.com

ABSTRACT

A number of animals with genetically identical appearance can be produced by somatic cell nuclear transfer (SCNT). From current advancement of SCNT and molecular techniques, production of a transgenic animal becomes easier. Although cloning efficiency in goat is low, the ability to propagate genetically identical animals, with a gene or genes of interest, would be important for increasing productivity and ultimately the economic livelihood. In this paper, the potential applications and uses of SCNT technology like production of transgenic goat for production of quality milk and meat are discussed.

Keywords:

Somatic Cell Nuclear Transfer, Goat, Applications

Received : 06 March 2014,

Revised: 30 March 2014,

Accepted : 06 April 2014,

Published online: 27 April 2014.

INTRODUCTION

Animal cloning is a potential technology for the production of genetically identical animals. Wilmut et al. (1997) could produce the first cloned sheep, namely "Dolly" by the transfer of a nucleus from a differentiated somatic cell into enucleated oocyte by the process of nuclear transfer (NT). Now, SCNT technology is expected to be useful for farm animal breeding and research, production of transgenic animals for biomedical purposes, and the conservation of endangered species. This technology needs more improvement due to the high rate of loss occurred during gestation and the abnormalities observed in

cloned animals (Constant et al., 2006; Heyman et al., 2002; Hoffert et al., 2005; Oishi et al., 2006). The milk obtained from the SCNT derived transgenic goat could produce antithrombin III (Baguisi et al., 1999). Thus, goats are important for the transgenic production of therapeutic recombinant proteins because of their high yield of purified product, relatively shorter generation interval and low incidence of the disease (Meade et al., 1998). Up to 700 liters of milk per year can be obtained from a single transgenic goat, capable of expressing between 1 and 10 g of protein per liter. Although milk yields are higher in dairy cows, dairy goats offer a shorter generation interval and thus better-looking for use in competitive transgenic programs (Ziomek, 1998). The production of transgenic animal derived from SCNT technology is more efficient method than pronuclear DNA-microinjection (Schnieke et al., 1997; Cibelli et al., 1998a; Ziomek, 1998; McCreath et al., 2000).

Transfected cells with gene of interest can be screened prior to nuclear transfer (NT) to detect both the presence and location of the transgene in the donor cell DNA. In addition, the advantage of this approach is that 100% of the resulting offspring will be transgenic (Niemann and Kues, 2000). Some of the proteins currently being produced in transgenic animals via SCNT are human clotting factor IX (Schnieke et al., 1997) and antithrombin III (Baguisi et al., 1999). The advantage of SCNT techniques in dairy goats to generate the human recombinant proteins and monoclonal antibodies in the milk of transgenic offspring could potentially contribute to the pharmaceutical industry (Baguisi et al., 1999). In this article, SCNT technology in goat has been discussed.

WHAT IS SCNT

Somatic cell nuclear transfer is a complex technique, involving donor cell selection, oocyte collection and enucleation, nuclear transfer, and embryo transfer. The most cells in normal mammalian development are gradually losing their pluripotency when develop into differentiated cells. The critical aspect of SCNT is reprogramming of differentiated cells, so the reconstructed donor cells and recipient cytoplasm can develop into live clones. Therefore, type of donor cell and quality of receptor cytoplasm affect efficiency of SCNT.

DONOR CELLS

Cloned goat have been generated from several cell types including cumulus cells (Zou et al., 2002; Keefer et al., 2002), fetal fibroblasts (Baguisi et al., 1999; Keefer et al., 2002; Melican et al., 2005; Liu et al., 2011; An et al., 2012; Soh et al., 2012; Liu et al., 2013), adult fibroblasts (Dutta et al., 2011), lymphocyte cells (Dutta et al., 2011), bone marrow-derived mesenchymal stem cells (Kwong et al., 2014), ear fibroblast cells (Wang et al., 2007; Abdullah et al., 2011; Liu et al., 2012; Kwong et al., 2012; Wan et al., 2012; Kwong et al., 2014), mammary gland epithelial Cells (Zhang et al., 2008; Yuan et al., 2009; Zhang et al., 2010; Yuan et al., 2014). However, the cell cycle of the donor cells usually affects reprogramming of the reconstructed embryos and their development. The G0 phase is a quiescent stage of cell cycle and after fusion into oocyte cytoplasm, the G0 nuclei of the oocytes were synchronized for DNA replication. The possibility of chromosomal aberrations during development of nuclear-transplanted cells could help cells to reprogram the normal cell cycle. In previous SCNT experiments, serum starvation, contact inhibition (Song et al., 2009; Jang et al., 2005), or chemical treatment (Gibbons et al., 2002; Yu et al., 2003; Vacková et al., 2003), were generally used to induce the cells into the G0/G1 phase. Melican et al. (2005) suggested that the kidding rate of the serum starvation treatment group was greater than contact inhibition treatment, indicating that the serum starvation treatment was more beneficial than the contact inhibition treatment.

RECIPIENT OOCYTES

Most current SCNT studies use MII-phase enucleated oocytes as recipient cytoplasm for NT. There are two sources for these stage oocytes: *in vivo* mature oocytes collected directly from animals or aspirated oocytes collected from the ovaries of live or slaughtered

animals and then cultured *in vitro* for maturation. In sheep, the different oocyte sources did not affect the fusion of reconstructed embryos, but blastocyst formation, recipient sheep pregnancy, and lambing rates of *in vivo* matured oocytes were significantly higher than those of *in vitro* matured oocytes (Well et al., 1997). However, in goats, there was no significant difference in developmental capabilities of *in vivo* and *in vitro* matured oocytes as recipient cytoplasm (Reggio et al., 2001). Thus, the most recipient oocytes for SCNT in goats are derived from slaughter house.

APPLICATION OF SCNT IN GOATS

SCNT is a relatively simple and inexpensive technique for increasing production of commercial value in dairy goat by using transgenic goats for the production of many recombinant therapeutic proteins for treatment of diseases. In fact, the first drug called ATryn, a human antithrombin protein approved for human use was derived from a genetically engineered goat (Dutta et al., 2011). Thus, goat is also an adapted domestic species for biological investigation and application, because of its short gestation period and easy maintenance. Moreover, SCNT technique was greatly improved, and widely successful in other animal species. Previously, the technology to produce transgenic goat was the microinjection of the DNA construct into the pronuclear embryos. However, this method shows a low efficiency and cannot predict outcome (Baldassarre and Karatzas, 2004). For SCNT, donor cell can be genetically modified by introducing the gene of interest and screened before use in SCNT.

SCNT in goats for milk production: Cloned goats express many proteins in their milk that cannot be chemically synthesized or manufactured by bacterial or cell culture systems (Baguisi et al., 1999; Reggio et al., 2001). Thus, the current progress of the SCNT technology has successfully produced healthy transgenic goats for the production of α -fetoprotein (AFP), suggesting that purified AFP derived from the milk of SCNT goat was similar to human AFP isolated from human cord blood (Parker et al., 2004). The milk of the four transgenic goats developed by SCNT could produce MSP142-malaria antigen (Behboodi et al., 2005). Recombinant lysosomal human acid β -glucosidase, the enzyme involves for catabolism of glucosylceramide into ceramide and glucose, has been successfully expressed in milk of cloned transgenic goat (Zhang et al., 2010). Human lactoferrin (hLTF), one of the components of the immune system of the body that provides antibacterial activity to human infants, can be produced from milk of SCNT goat (An et al., 2012; Wan

et al., 2012; Meng et al., 2013). Milk from cloned transgenic goat produced by SCNT could express human lysozyme (*hLYZ*), which possesses antimicrobial properties causing increase in safety and storage time of milk (Liu et al., 2013). Moreover, Yekta et al. (2013) showed great success in SCNT transgenic goat using fetal fibroblasts with a linearized marker-free construct in which the transgene was juxtaposed to b-casein promoter designed to secrete the recombinant protein in goat milk. Greater expression of recombinant human alpha-lactalbumin derived from transgenic goat using SCNT was observed in goat milk (Yuan et al., 2014).

SCNT in goats for meat production: Goat meat is an important food worldwide and it has high-quality protein and low cholesterol; thus providing health benefits to humans. SCNT using myostatin-targeted 2-month-old goat fibroblasts as donor is a potential method for producing myostatin-targeted goats for meat purpose (Zhou et al., 2013).

CONCLUSIONS

The ability of SCNT technology to propagate genetically identical animals with superior genotypes would be important for increasing the number of economically important animals. Importantly, this technology can be applied with cell culture system and animal molecular technology to produce transgenic animal. However, the efficiency of animal production derived from SCNT is still low. Moreover, the production of animals by this technology could lead to more efficient models for biomedical research.

REFERENCES

- Abdullah RB, Wan Khadijah WE, Kwong PJ (2011). Comparison of intra- and interspecies nuclear transfer techniques in the production of cloned caprine embryos. *Small Ruminant Research*, 98: 196-200.
- An LY, Yuan YG, Yu BL, Yang TJ, Cheng Y (2012). Generation of human lactoferrin transgenic cloned goats using donor cells with dual markers and a modified selection procedure. *Theriogenology*, 78: 1303-1311.
- Baguisi A, Behboodi E, Melican DT, Pollock JS, Destrempes MM, Cammuso C, Williams JL, Nims SD, Porter CA, Midura P, Palacios MJ, Ayres SL, Denniston RS, Hayes ML, Ziomek CA, Meade HM, Godke RA, Gavin WG, Overström EW (1999). Production of goats by somatic cell nuclear transfer. *Nature Biotechnology*, 17: 456-461.
- Baldassarre H, Karatzas CN (2004). Advanced assisted reproduction technologies (ART) in goats. *Animal Reproduction Science*, 82-83: 255-266.
- Behboodi E, Ayres SL, Memili E, O'Coin M, Chen LH, Reggio BC, Landry AM, Gavin WG, Meade HM, Godke RA, Echelard Y (2005). Health and reproductive profiles of malaria antigen-producing transgenic goats derived by somatic cell nuclear transfer. *Cloning and Stem Cells*, 7: 107-118.
- Cibelli JB, Stice SL, Golueke PJ, Kane JJ, Jerry J, Blackwell C, Ponce de León FA, Robl JM (1998a). Cloned transgenic calves produced from nonquiescent fetal fibroblasts. *Science*, 280: 1256-1258.
- Constant F, Guillomot M, Heyman Y, Vignon X, Laigre P, Servely JL, Renard JP, Chavatte-Palmer P (2006). Large offspring or large placenta syndrome? Morphometric analysis of late gestation bovine placentomes from somatic nuclear transfer pregnancies complicated by hydrallantois. *Biology of Reproduction*, 75: 122-130.
- Dutta R, Malakar D, Khate K, Sahu S, Akshey Y, Mukesh M (2011). A comparative study on efficiency of adult fibroblast, putative embryonic stem cell and lymphocyte as donor cells for production of handmade cloned embryos in goat and characterization of putative ntES cells obtained from these embryos. *Theriogenology*, 76: 851-863.
- Gibbons J, Arat S, Rzucidlo J, Miyoshi K, Waltenburg R, Respass D (2002). Enhanced survivability of cloned calves derived from roscovitine-treated adult somatic cells. *Biology of Reproduction*, 66: 895-900.
- Heyman Y, Chavatte-Palmer P, LeBourhis D, Camous S, Vignon X, Renard JP (2002). Frequency and occurrence of late-gestation losses from cattle cloned embryos. *Biology of Reproduction*, 66: 6-13.
- Hoffert KA, Batchelder CA, Bertolini M, Moyer AL, Famula TR, Anderson DL, Anderson GB (2005). Measures of maternal-fetal interaction in day-30 bovine pregnancies derived from nuclear transfer. *Cloning and Stem Cells*, 7: 289-305.
- Jang G, Jeon HY, Ko KH, Park HJ, Kang SK, Lee BC (2005). Developmental competence and gene expression in preimplantation bovine embryos derived from somatic cell nuclear transfer using different donor cells. *Zygote*, 13: 187-95.
- Keefer CL, Keyston R, Lazaris A, Bhatia B, Begin I (2002). Production of cloned goats after nuclear transfer using adult somatic cells. *Biology of Reproduction*, 66: 199-203.

- Kwong PJ, Abdullah RB, Wan Khadijah WE (2012). Increasing glucose in KSOMaa basal medium on culture Day 2 improves *in vitro* development of cloned caprine blastocysts produced via intraspecies and interspecies somatic cell nuclear transfer. *Theriogenology*, 78: 921–929.
- Kwong P, Nam H, Wan Khadijah W, Kamarul T, Abdullah R (2014). Comparison of *in vitro* developmental competence of cloned caprine embryos using donor karyoplasts from adult bone marrow mesenchymal stem cells vs ear fibroblast cells. *Reproduction in Domestic Animal*, doi: 10.1111/rda.12262.
- Liu J, Li LL, Du S, Bai XY, Zhang HD (2011). Effects of interval between fusion and activation, cytochalasin B treatment, and number of transferred embryos, on cloning efficiency in goats. *Theriogenology*, 76: 1076–1083.
- Liu J, Luo Y, Liu Q, Zheng L, Yang Z, Wang Y, Su J, Quan F, Zhang Y (2012). Production of cloned embryos from caprine mammary epithelial cells expressing recombinant human beta-defensin-3. *Theriogenology*, 79: 660–666.
- Liu J, Luo Y, Ge H, Han C, Zhang H, Wang Y, Su J, Quan F, Gao M, Zhang Y (2013). Anti-bacterial activity of recombinant human *b*-defensin-3 secreted in the milk of transgenic goats produced by somatic cell nuclear transfer. *PLoS ONE*, 8: e65379–e65379.
- McCreath KJ, Howcroft J, Campbell KH, Colman A, Schnieke AE, Kind AJ (2000). Production of gene-targeted sheep by nuclear transfer from cultured somatic cells. *Nature*, 405: 1066–1069.
- Meade HM, Echelard Y, Ziomek CA, Young MW, Harvey M, Cole ES, Groet S, Smith TE, Curling JM (1998). Expression of recombinant proteins in the milk of transgenic animals (Fernandez JM, Hoeffler JP, Eds). In: *Gene Expression Systems: Using Nature for the Art of Expression*; pp 399–427.
- Melican D, Butler R, Hawkins N, Chen LH, Hayden E, Destrempe M (2005). Effect of serum concentration, method of trypsinization and fusion/activation utilizing transfected fetal cells to generate transgenic dairy goats by somatic cell nuclear transfer. *Theriogenology*, 63: 1549–1463.
- Meng L, Jia RX, Sun YY, Wang ZY, Wan YJ, Zhang YL, Zhong BS, Wang F (2013). Growth regulation, imprinting and epigenetic transcription-related gene expression differs in lung of deceased transgenic cloned and normal goats. *Theriogenology*, 81: 459–466.
- Niemann H, Kues WS (2000). Transgenic livestock: Premises and promises. *Animal Reproduction Science*, 60: 277–293.
- Oishi M, Gohma H, Hashizume K, Taniguchi Y, Yasue H, Takahashi S, Yamada T, Sasaki Y (2006). Early embryonic death-associated changes in genome-wide gene expression profiles in the fetal placenta of the cow carrying somatic nuclear derived cloned embryo. *Molecular Reproduction and Development*, 73: 404–409.
- Parker MH, Birck-Wilson E, Allard G, Masiello N, Day M, Murphy KP, Paragas V, Silver S, Moody MD (2004). Purification and characterization of a recombinant version of human alpha-fetoprotein expressed in the milk of transgenic goats. *Protein Expression and Purification*, 38: 177–183.
- Reggio BC, James AN, Green HL, Gavin WG, Behboodi E, Echelard Y (2001). Cloned transgenic offspring resulting from somatic cell nuclear transfer in the goat: oocytes derived from both follicle-stimulating hormone-stimulated and nonstimulated abattoir-derived ovaries. *Biology of Reproduction*, 65: 1528–1533.
- Schnieke AE, Kind AJ, Ritchie WA, Mycock K, Scott AR, Ritchie M, Wilmut I, Colman A, Campbell KH (1997). Human factor IX transgenic sheep produced by transfer of nuclei from transfected fetal fibroblasts. *Science*, 278: 2130–2133.
- Soh HH, Wan Khadijah WE, Abdullah RB (2012). Subzonal versus intracytoplasmic injection produces a higher rate of cloned caprine-bovine interspecies blastocysts. *Small Ruminant Research*, 105: 231–236.
- Song K, Hyun SH, Shin T, Lee E (2009). Post-activation treatment with demecolcine improves development of somatic cell nuclear transfer embryos in pigs by modifying the remodeling of donor nuclei. *Molecular Reproduction and Development*, 76: 611–9.
- Vacková I, Engelová M, Marinov I, Tománek M (2003). Cell cycle synchronization of porcine granulosa cells in G1 stage with mimosine. *Animal Reproduction Science*, 77: 235–245.
- Wan YJ, Zhang YL, Zhou ZR, Jia RX, Li M, Song H, Wang ZY, Wang LZ, Zhang GM, You JH, Wang F (2012). Efficiency of donor cell preparation and recipient oocyte source for production of transgenic cloned dairy goats harboring human lactoferrin. *Theriogenology*, 78: 583–592.
- Wang L, Peng T, Zhu H, Lv Z, Liu TT, Shuai ZQ, Gao H, Cai T, Cao X, Wang HQ (2007). *In vitro* development of reconstructed ibex (*Capra ibex*) embryos by nuclear transfer using goat (*Capra*

- hircus*) oocytes. *Small Ruminant Research*, 73: 135-141.
- Wells DN, Misica PM, Day TA, Tervit HR (1997). Production of cloned lambs from an established embryonic cell line: a comparison between in vivo- and in vitro-matured cytoplasts. *Biology of Reproduction*, 57: 385-393.
- Wilmot I, Schnieke AE, McWhir J, Kind AJ, Campbell KH (1997). Viable offspring derived from fetal and adult mammalian cells. *Nature*, 385: 810-813.
- Yekta AA, Dalman A, Eftekhari-Yazdi P, Sanati MH, Shahverdi AH, Fakheri R, Vazirinasab H, Daneshzadeh MT, Vojgani M, Zomorodipour A, Fatemi N, Vahabi Z, Mirshahvaladi S, Ataei F, Bahraminejad E, Masoudi N, Valojerdi MR, Gourabi H (2013). Production of transgenic goats expressing human coagulation factor IX in the mammary glands after nuclear transfer using transfected fetal fibroblast cells. *Transgenic Research*, 22: 131-142.
- Yu YS, Sun XS, Jiang HN, Han Y, Zhao CB, Tan JH (2003). Studies of the cell cycle of in vitro cultured skin fibroblasts in goats: work in progress. *Theriogenology*, 59: 1277-1289.
- Yuan YG, Cheng Y, Guo L, Ding GL, Bai YJ, Miao MX, An LY, Zhao JH, Cao YJ (2009). Cloned kids derived from caprine mammary gland epithelial cells. *Theriogenology*, 72: 500-505.
- Yuan YG, An L, Yu B, Song S, Zhou F, Zhang L, Gu Y, Yu M, Cheng Y (2014). Expression of Recombinant Human Alpha-Lactalbumin in the Milk of Transgenic Goats Using a Hybrid Promoter/Enhancer, doi: 10.1155/2014/281031.
- Zhang J, Li L, Cai Y, Xu X, Chen J, Wu Y (2008). Expression of active recombinant human lactoferrin in the milk of transgenic goats. *Protein Expression and Purification*, 57: 127-35.
- Zhang YL, Wan YJ, Wang ZY, Xu D, Pang XS, Meng L (2010). Production of dairy goat embryos, by nuclear transfer, transgenic for human acid beta-glucosidase. *Theriogenology*, 73: 681-90.
- Zhou ZR, Zhong BS, Jia RX, Wan YJ, Zhang YL, Fan YX, Wang LZ, You JH, Wang ZY, Wang F (2013). Production of myostatin-targeted goat by nuclear transfer from cultured adult somatic cells. *Theriogenology*, 79: 225-33.
- Ziomek CA (1998). Commercialization of proteins produced in the mammary gland. *Theriogenology*, 49: 139-144.
- Zou X, Wang Y, Cheng Y, Yang Y, Ju H, Tang H (2002). Generation of cloned goats (*Capra hircus*) from transfected foetal fibroblast cells, the effect of donor cell cycle. *Molecular Reproduction and Development*, 61: 164-172.