

ORIGINAL RESEARCH

Effect of aromatic oils on the expression of some virulence-associated and antimicrobial resistance genes of *Escherichia coli* isolated from broilers

Walid Hamdy Hassan¹ , Hala Sayed Hassan Salam¹ , Wafaa Mohamed Hassan² , Salama Abohamra Sayed Shany³ , Ghada Sayed Ibrahim Osman⁴ 

¹Department of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef, Egypt

²Microbiology-Reference Laboratory of Quality Control on Poultry Production, Animal Health Research Institute AHRI, Giza Governorate, Egypt

³Department of Poultry Diseases, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef, Egypt

⁴Department of Bacteriology, Animal Health Research Institute, Beni-Suef Laboratory, Agricultural Research Center, Beni-Suef, Egypt

ABSTRACT

Objectives: This study aimed to prove the effects of *Escherichia coli* isolates isolated from diseased broilers to form biofilms, describe their antimicrobial sensitivity, and determine the effect of allicin and cinnamon essential oils on the expression of some genes (*fimH*, *int1*, and *luxS*) through quantitative polymerase chain reaction (q-PCR).

Materials and Methods: 140 samples were obtained from diseased broilers in Beni-Suef Governorate, Egypt. These samples were examined by conventional bacteriology methods to detect the causative agent. The antimicrobial susceptibility of the isolated bacteria was assessed using the disc diffusion method, The ability of yeast extract-casamino acids Congo Red Agar to generate phenotypic biofilms was next tested. The presence of resistance and virulence genes in some multidrug resistant isolates was genotypically investigated. The antibacterial effects of allicin and cinnamon oil were evaluated against the growth of multidrug-resistant *E. coli*. Finally, q-PCR was utilized to assess changes in some genes' expression.

Results: *Escherichia coli* was isolated from 61 samples (43.6%). An antimicrobial susceptibility test revealed that multidrug-resistance (MDR) (could resist more than three antimicrobial classes) *E. coli* prevalence was 100%. 40.8% of isolates phenotypically produce biofilms. The detection of resistance and virulence genes by PCR showed that all tested isolates carry *aadB*, *fimH*, *int1*, *qnrS*, and *luxS* genes, while only 40% harbor *iss* genes. q-PCR showed that after treatment with allicin and cinnamon oils, gene expression went down.

Conclusion: This investigation highlights that *E. coli* showed resistance against most of the tested antimicrobials; all isolates were MDR. The study showed wide dissemination of virulence and resistance genes among *E. coli*. Allicin and cinnamon oils have antimicrobial activities and could be used as alternatives to synthetic antimicrobial agents.

ARTICLE HISTORY

Received August 01, 2021

Revised September 08, 2021

Accepted September 16, 2021

Published June 26, 2022

KEYWORDS

Escherichia coli; broiler; virulence; resistance; aromatic oils



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

Introduction

Escherichia coli belongs to the Enterobacteriaceae family [1]. Even though *E. coli* is a normal occupant of poultry's digestive tract, and pathogenic serotypes have previously been reported from the intestinal tract of healthy poultry, supporting the claim that *E. coli* is frequently a secondary or opportunistic infection [2]. *Escherichia coli* can cause

colibacillosis outbreaks associated with various disease conditions [3]. Stress may play a role in transmitting avian pathogenic *E. coli* from the bird's gut to the bloodstream. It infects several internal organs and produces pericarditis, perihepatitis, salpingitis, peritonitis, and other extra-intestinal symptoms [4]. Colibacillosis can affect chickens of different ages but is more common and severely affects

Correspondence Hala Sayed Hassan Salam ✉ hala_saydh@yahoo.com 📧 Department of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef, Egypt.

How to cite: Hassan WH, Salam HSH, Hassan WM, Shany SAS, Osman GSI. Effect of aromatic oils on the expression of some virulence-associated and antimicrobial resistance genes of *Escherichia coli* isolated from broilers. J Adv Vet Anim Res 2022; 9(2):191–202.

young birds [2]. Colisepticemia is the most prevalent form of avian colibacillosis, which causes hefty financial losses worldwide [5]. Misusage of antimicrobials and their prophylactic application in the poultry field are considered to be the causes of drug resistance [6]. Increased antimicrobial resistance patterns pose an important public health concern [7].

Therefore, the drug of choice should be chosen based on its sensitivity, which may be determined by lab examination and employed for an accurate dose and course. Other than resistance gene expression, bacteria can protect themselves by producing vast amounts of extracellular polymeric substance (EPS) during the biofilm formation process [8]. This EPS plays a critical role in antibiotic resistance and host immunity [9]. Quorum sensing (QS), a type of cell-to-cell communication in which an accumulation of signaling molecules in the extracellular environment causes gene expression to be controlled, initiates the biofilm-building process [10]. Several spices and plants are now prized for their antibacterial properties [11]. Consequently, there is an increased interest in using natural, non-toxic, and effective compounds as an alternative to synthetic antimicrobials. These are primarily herb and spice extracts and essential oils (EOs) [12]. The Eos antimicrobial activity might be due to the high hydrophobicity that enables them to cross the microbial cell membranes, leading to loss of function or damage of the cellular proteins, lipids, and organelles, resulting in bacterial cell death [13]. The cinnamon EO's antibacterial activity may be attributed to its richness in cinnamaldehyde, which exhibits antibacterial properties against many pathogenic bacteria [14]. Garlic contains diallyl sulfide, which penetrates and dissolves the bacteria's protective bio-layer, eventually killing them [15]. Allicin, a component of garlic, contains a variety of thiosulfonates responsible for its antibacterial properties [16]. The goal of this study was to find out if *E. coli* from sick broilers in Beni-Suef Governorate, Egypt, had the ability to form biofilms and if they were sensitive to antimicrobials. They also wanted to use quantitative polymerase chain reaction (q-PCR) to find out how allicin and cinnamon essential oils affect the expression of certain genes (*fimH*, *int1*, and *luxS*).

Materials and Methods

Samples

One hundred forty samples were taken aseptically from broiler chicks aged one to 35 days and collected from Beni-Suef governorate, Egypt, between January 2020 and April 2020. These chicks suffered from respiratory manifestations and/or gastrointestinal problems. Swabs from the heart, lung, liver, and yolk sac that showed gross lesions were used for bacteriological isolation.

Bacteriological examination

Bacteriological and biochemical identification of isolated *E. coli*

The samples taken were inoculated under aseptic conditions into buffered peptone water and incubated aerobically at 37°C for 24 h. A loopful from the media was streaked onto MacConkey agar medium. Then all plates were incubated aerobically at 37°C for 24 h. Suspicious colonies for *E. coli* appear on MacConkey agar as pinkish colonies. One colony representing a single typical colonial appearance and morphological character was picked up for inoculation onto eosin methylene blue agar medium. Bacterial smears were stained by Gram's staining technique. Biochemical identification was carried out according to Quinn et al. [17], including oxidase test, indole production, methyl red test, Voges Proskauer test, citrate utilization test, sugar fermentation test, and hydrogen sulfide (H₂S) production on TSI medium, and urea hydrolysis test.

Antimicrobial sensitivity testing of *E. coli* isolates

The antibiogram was done by using the disk diffusion method. Antimicrobial discs included amoxicillin/clavulanic acid (30 µg), cefotaxime (30 µg), doxycycline (30 µg), fosfomycin (200 µg), gentamicin (10 µg), nitrofurantoin (300 µg), norfloxacin (10 µg), oxytetracycline (30 µg) and sulphamethoxazole/trimethoprim (25 µg), represented different antimicrobial classes of veterinary concern. The recorded zones were interpreted according to CLSI [18]. Multidrug-resistance (MDR) was defined as resistance to three or more antimicrobial classes according to Magiorakos et al. [19].

Biofilm formation of *E. coli* isolates

The test was done according to Zhou et al. [20]. YESCA/CR agar plates were inoculated and incubated aerobically for 48 h at 26°C. Positive results were indicated by dark red, red, or light red colonies, while negative results were pink, light pink, or white colonies.

Detection virulence-associated and resistance genes in *E. coli*

Five MDR *E. coli* isolates were subjected to PCR to identify three resistance-associated genes (*aadB*, *int1* and *qnrS*) and three virulence-related genes (*fimH*, *iss*, and *luxS*). The QIAamp DNA mini kit was used to extract DNA, as stated by the manufacturer's instructions and using their specific forward and reverse primers as indicated in Table 1. The DNA of the positive control was taken from an *E. coli* field isolate that was proven to be positive in RLQP (Reference Laboratory for Veterinary Quality Control on Poultry

Production, Dokki, Giza, Egypt). A negative control, on the other hand, is a PCR mixture without the DNA template.

Evaluation of the antibacterial effects of allicin and cinnamon oils on the growth of multidrug-resistant *E. coli* isolates

Allicin and cinnamon oils were investigated for their antimicrobial activities against 15 MDR *E. coli* isolates [21]. Essential oils were prepared as 10% (10,000 ppm) in DMSO as the original solution. Allicin oil (diallyl disulfide, 99.9%) was obtained from Seven Seas Pure Chemical Industries, Ltd., China. It was prepared at 500, 250, and 125 ppm concentrations in tryptone soya agar. Cinnamon oil (Aldrick, Sigma) was prepared at 125, 63, 32, 16, and 8 ppm concentrations in tryptone soya agar.

Qualitative real-time PCR for *E. coli* isolates

Three MDR *E. coli* strains were subjected to quantitative RT-PCR before and after treatment with 125 ppm allicin

and 32 ppm cinnamon to estimate variations of *fimH* luxS, and *Int1* genes. A quantitative RT-PCR was prepared using the Quantitect SYBR Green PCR kit. Primer sequences and probes used in SYBR Green real-time PCR are illustrated in Table 2.

Results

Prevalence of *E. coli*

Sixty-one *E. coli* isolates were isolated from 140 samples collected from diseased broilers with an incidence of 43.6 %.

Antimicrobial sensitivity testing of recovered *E. coli*

The explored *E. coli* isolates ($n = 61$) showed high sensitivity rates to fosfomycin (85.2%) and nitrofurantoin (80.3%). On the contrary, high resistance rates were recorded for amoxicillin-clavulanic acid (100%), cefotaxime (98.4%), oxytetracycline (90.2%), sulfatrimethoprim (90.2%), and

Table 1. Primers of virulence-associated and resistance genes used in PCR.

Target gene	Primers direction	Primers sequences (5'-3')	Amplified product size (bp)	Reference
<i>AadB</i>	F	GAG CGA AAT CTG CCG CTC TGG	319	[61]
	R	CTG TTA CAA CGG ACT GGC CGC		
<i>FimH</i>	F	TGC AGA ACG GAT AAG CCG TGG	508	[62]
	R	GCA GTC ACC TGC CCT CCG GTA		
<i>lss</i>	F	ATG TTA TTT TCT GCC GCT CTG	266	[63]
	R	CTA TTG TGA GCA ATA TAC CC		
<i>Int1</i>	F	CCT CCC GCA CGA TGA TC	280	[35]
	R	TCC ACG CAT CGT CAG GC		
<i>LuxS</i>	F	ATG CCG TTG TTA GAT AGC TTC A	513	[64]
	R	GAT GTG CAG TTC CTG CAA CTT C		
<i>QnrS</i>	F	ACG ACA TTC GTC AAC TGC AA	417	[65]
	R	TAA ATT GGC ACC CTG TAG GC		

Table 2. Oligonucleotide primers and probes used in SYBR Green real time PCR.

Gene	Primers direction	Primer sequence (5'-3')	Reference
<i>E. coli</i> 16S rRNA	F	GCT GAC GAG TGG CGG ACG GG	[66]
	R	TAG GAG TCT GGA CCG TGT CT	
<i>FimH</i>	F	TGC AGA ACG GAT AAG CCG TGG	[62]
	R	GCA GTC ACC TGC CCT CCG GTA	
<i>LuxS</i>	F	ATG CCG TTG TTA GAT AGC TTC A	[64]
	R	GAT GTG CAG TTC CTG CAA CTT C	
<i>Int1</i>	F	CCT CCC GCA CGA TGA TC	[35]
	R	TCC ACG CAT CGT CAG GC	

gentamicin (88.5%). The detailed results of antimicrobials are included in Table 3. Moreover, all the investigated *E. coli* isolates (100%) were MDR.

Biofilm formation of *E. coli* on YESCA/CR agar

Out of 61 *E. coli* isolates, 37 (60.7%) were positive on YESCA/CR agar, while 24 were negative on YESCA/CR agar.

PCR of *E. coli*

All of the tested *E. coli* isolates ($n = 5$) carried *aadB*, *int1* and *qnrS* genes (100%) in terms of the resistance genes.

In relation to the virulence-associated genes, all the tested *E. coli* isolates (100%) had *fimH*, and *luxS*; meanwhile, two tested *E. coli* isolates (40%) harbored *iss* gene. (Figs. 1–3).

Antibacterial effects of allicin and cinnamon oils on multi-drug-resistant *E. coli* isolates

Allicin oil fully inhibited the growth of all *E. coli* isolates ($n = 15$) (100%) at concentrations of 250 and 500 ppm. On the contrary, 125 and 63 ppm of allicin concentrations lead to retarded and little growth of *E. coli* (Table 4).

Cinnamon oil fully inhibited the growth of all *E. coli* isolates ($n = 15$) (100%) at concentrations of 125 ppm and 63

Table 3. Antimicrobial susceptibility of *E. coli* isolated from broiler chicks ($n = 61$).

Antimicrobial class	Antimicrobial disk	Susceptible		Intermediate		Resistant	
		No.	%	No.	%	No.	%
β -lactamase stable	Amoxicillin-clavulanic acid	0	0	0	0	61	100
Cephalosporins	Cefotaxime	0	0	1	1.6	60	98.4
Tetracyclines	Doxycycline	7	11.5	7	11.5	47	77
	Oxytetracycline	2	3.3	4	6.5	55	90.2
Fosfomycins	Fosfomicin	52	85.2	0	0	9	14.8
Aminoglycosides	Gentamicin	3	4.9	4	6.6	54	88.5
Nitrofurans	Nitrofurantoin	49	80.3	10	16.4	2	3.3
Quinolones	Norofloxacin	3	4.91	11	18.03	47	77.04
Potentiated sulfonamide	Sulphamethoxazole/trimethoprim	3	4.9	3	4.9	55	90.2




Figure 1. PCR amplification of the *int1* gene at 280bp and the *qnrS* gene at 417bp fragments. Lanes 1–5 showed positive amplification of *int1* and *qnrS* genes. P = Positive control; N = negative control; L = 100-bp DNA molecular size ladder.




Figure 2. PCR amplification of the *iss* gene at 266 bp and the *fimH* gene at 508 bp fragments. Lanes 1–5 showed positive amplification of *fimH* gene, and Lanes 3, 5 showed positive amplification of *iss* gene. P = Positive control; N = negative control; L = 100-bp DNA molecular size ladder.




Figure 3. PCR amplification of the *aadB* gene at 319 bp and the *luxS* gene at 513 bp fragments. Lanes 1–5 showed positive amplification of *aadB* and *luxS* genes. P = Positive control; N = negative control; L = 100-bp DNA molecular size ladder.

ppm, while at 32 ppm it inhibited 4 isolates only (26.7%) of *E. coli*. On the other hand, the 16 and 8 ppm concentrations did not kill the *E. coli* isolates, but they grew slowly and not at all (Table 5).

Quantitative real time-PCR (RT-PCR) for multidrug-resistant *E. coli* before and after treatment with allicin and cinnamon oils

Table 6 and Figures 4–7 illustrated the down-regulation in fold change of *E. coli* for the *fimH* gene after exposure

to 125 ppm allicin and 32 ppm cinnamon oil were 0.4061 and 0.1397, respectively, for the first *E. coli* isolate; 0.4118 and 0.0693 for the second *E. coli* isolate; and 0.3763 and 0.1768, respectively, for the third *E. coli* isolate. In the same context, down regulations in fold change for *Int1* gene were 0.6974 and 0.3099 for the first *E. coli* isolate, 0.6417 and 0.2755, respectively, for the second *E. coli* isolate, and 0.5105 and 0.2755 for the third *E. coli* isolate. On the other hand, down regulations in fold change for *luxS* were 0.7120 and 0.3121, respectively, for the first *E. coli* isolate; 0.6156

Table 4. The antibacterial effect of allicin oil on *E. coli* isolates (*n* = 15).

Bacterial isolates	Allicin oil concentration							
	500 ppm		250 ppm		125 ppm		63 ppm	
	No.	%	No.	%	No.	%	No.	%
<i>E. coli</i>	0	0	0	0	15	100	15	100

No.: Number of the affected (growth inhibited) isolates.

%: was calculated according to the number (No.) of tested bacterial isolates.

Table 5. The antibacterial effect of cinnamon oil on *E. coli* isolates (*n* = 15).

Bacterial isolates	Cinnamon oil concentration									
	8 ppm		16 ppm		32 ppm.		63 ppm		125 ppm	
	No.	%	No.	%	No.	%	No.	%	No.	%
<i>E. coli</i>	0	0	0	0	4	26.7	15	100	15	100

No.: Number of the affected (growth inhibited) isolates.

%: was calculated according to the number (No.) of tested bacterial isolates.

Table 6. RT-PCR analysis for *E. coli* isolates before and after treatment Allicin and cinnamon oils.

<i>E. coli</i> Sample No.	Treatment	Sample ID	16S rRNA		fimH		luxS		Int1	
			CT	CT	Fold change	CT	Fold change	CT	Fold change	
1	Control	A1	19.87	20.38	-	23.06	-	21.46	-	
	Allicin	B1	20.52	22.33	0.4061	24.20	0.7120	22.63	0.6974	
	Cinnamon	C1	19.86	23.21	0.1397	24.73	0.3121	23.14	0.3099	
2	Control	A2	20.11	20.92	-	22.95	-	22.18	-	
	Allicin	B2	19.44	21.53	0.4118	22.98	0.6156	22.15	0.6417	
	Cinnamon	C2	18.91	23.57	0.0693	24.12	0.1934	22.84	0.2755	
3	Control	A3	20.34	21.60	-	23.10	-	21.59	-	
	Allicin	B3	18.10	20.77	0.3763	21.13	0.8293	20.32	0.5105	
	Cinnamon	C3	19.62	23.38	0.1768	23.95	0.3368	22.73	0.2755	

and 0.1934, respectively, for the second *E. coli* isolate; and 0.8293 and 0.3368, respectively, for the third *E. coli* isolate.

Discussion

Escherichia coli infection causes severe economic losses in poultry production [22]. In this study, *E. coli* prevalence was studied in diseased broilers. Sixty-one isolates were recovered from the studied cases, with 43.6%. This finding was nearly identical to that reported by Awad et al. [23], who revealed only 28 *E. coli* isolates out of 54 (51.9%) in Egypt's northern delta during their first two weeks of life.

The disc diffusion method was used to study the *in vitro* susceptibility patterns of *E. coli* isolates against nine different antimicrobial drugs from eight other classes. The most

effective antimicrobials on *E. coli* were fosfomycin (85.2%) and nitrofurantoin (80.3%). The antimicrobial amoxicillin-clavulanic acid showed the highest resistance (100%), followed by cefotaxime (98.4%), oxytetracycline, sulfamethoxazole (90.2% for each), and gentamicin (88.5%).

The nearly same result of high resistance recorded by Amer et al. [24] oxytetracycline resistance was 85%. The reported high rates of resistance could be due to the use of antimicrobial drugs in a frequent manner [25]. *Escherichia coli* frequently carries many drug resistance plasmids and can easily transfer those plasmids to other species when stressed. *E. coli* can receive and transfer plasmids from one bacteria to another when there is a mixing of species in the intestines [26]. In this study, 100% of *E. coli* isolates were MDR. These findings corroborate those obtained by many




Figure 4. Down-regulation in *16s rRNA* gene from 3 MDR *E. coli* before and after exposure to 125 ppm allicin and 32 ppm cinnamon oils.




Figure 5. Down-regulation in *fimH* gene from 3 MDR *E. coli* before and after exposure to 125 ppm allicin and 32 ppm cinnamon oils.




Figure 6. Down regulation in *int1* gene from 3 MDR *E. coli* before and after exposure to 125 ppm allacin and 32 ppm cinnamon oils.




Figure 7. Down regulation in *luxS* gene from 3 MDR *E. coli* before and after exposure to 125 ppm allacin and 32 ppm cinnamon oils.

authors in Egypt who found that *E. coli* isolates were 100% MDR [7,24].

Biofilm formation is one of bacteria's resistance and virulence strategies [27], in which biofilms are surface-attached microbial cells encapsulated in an extracellular matrix that are much more resistant to antimicrobial treatments than non-adherent planktonic cells. Biofilm formation interferes with antimicrobial susceptibility. Bacteria become resistant via a variety of mechanisms, including antimicrobial interactions with biofilm matrix components, slower growth rates, and the various actions of specific antibiotic genetic determinants, which can increase antimicrobial resistance by up to 1,000 fold [28]. low doses of certain antibiotics can induce biofilm formation [27]. Reichhardt et al. [29] stated that Congo red on YESCA agar binds to curled whole cells without bacterial growth inhibition and can be used to measure whole-cell curling in which *E. coli* accumulates extracellular sticky amyloid fibers known as curli, which enables bacterial adherence and aids biofilm development.

In the present study, 60.7% of *E. coli* isolates were phenotypically positive biofilm formers on YESCA/CR agar. These results are higher than those of Amer et al. [24], 35.7%, and Raheel et al. [30], 46.6%, and lower than that obtained by Wang et al. [31], 81.64%.

A gentamicin resistance encoding gene, *aadB*, was identified in all tested *E. coli*. The high prevalence of the *aadB* gene in selected *E. coli* isolates was nearly similar to Zajaç et al. [32], which reported *aadB* in 93.8% of tested *E. coli* isolates. But lower percentages were obtained by Kim et al. [33] and Nhung et al. [34]: 2.5% and 12.3%, respectively, while Radwan et al. [12] did not detect the *aadB* gene in any isolates.

The antimicrobial resistance in APEC could be due to class 1 integron gene cassettes [33]. There is a strong relationship between the incidence of integrons and the phenotypes of multiresistance. Integrons carrying antimicrobial resistance genes have recently been more common in *E. coli* isolates and act as a vehicle for antimicrobial resistance acquisition and spread in the environment [35]. The *qnrS* gene is a plasmid-borne quinolone resistance gene. The *qnrS* gene of quinolones in *E. coli* isolates in this study poses a threat to public health. This is because *qnr*-plasmids are frequently coupled with integrons and contain a variety of resistance determinants, providing resistance to a variety of antimicrobial classes, including -lactams and aminoglycosides [36]. All tested *E. coli* isolates (100%) were positive for the *int* and *qnrS* genes. The high prevalence of the *int1* gene agrees with Ibrahim et al. [37] and is lower than Tran et al. [38], at 22.2%. The results of *qnrS* in the current study are lower than the result reported as 31.8% by Szmolka et al. [39].

The *fimH* gene is one of the adhesion encoding genes. Fim H is an APEC virulent factor that encodes a fimbrial

adhesion (type 1 pili) and promotes bacterial colonization, invasion, and biofilm formation [40]. The *fimH* gene was found in all tested isolates in the present study. The *fimH* results agreed with Wu et al. [41], who discovered that 100% of the tested isolates' genes were *fimH* positive.

Iss is a serum survival gene. This gene is significantly associated with APEC strains, and it is discovered in APEC significantly more frequently than commensal *E. coli* [42]. The presence of the *iss* gene in conjugative Col V plasmids suggests a link to APEC pathogenicity [43]. A similar result was reported by Moon et al. [44], who recorded *iss* as 41%.

In response to various environmental stimuli, *E. coli* uses QS to respond to their cell population and coordinate virulence gene expression [45]. Bacteria communicate via specialized chemical molecules to control the production of virulence factors and antimicrobial-resistant determinants to regulate their action against the host [46]. *luxS* (luciferase genes) has been found to have an essential function in regulating bacterial behavior [47]. It is also crucial for biofilm development, adhesion, and gene expression [31,48]. The LuxS/AI-2 QS system implicates the synthesis of cell signaling molecules via *luxS*-based autoinducer-2 (AI-2) [49]. The current work found the *luxS* gene in 100% of the tested isolates.

As a result of their anti-QS activity, some of these EOs demonstrate an inhibitory effect against resistant bacterial isolates, which could help lower the *in vivo* virulence and pathogenicity of drug-resistant bacteria [50]. The mode of action of the EOs depends on the cell wall degradation, cell membrane damage, cytoplasm coagulation, damaging the proteins of the membrane and increasing permeability, leading to cell content leakage (ions and other cell contents), decreasing the motive force of protons, decreasing the intracellular ATP pool due to decreased ATP synthesis and enhanced hydrolysis that is separate from the increased membrane permeability, and decreasing the membrane potential due to increased membrane permeability [50].

In this study, allicin showed complete growth inhibition of all the tested *E. coli* isolates (100%) at concentrations of 250 and 500 ppm. Allicin works primarily by blocking enzymes necessary for bacterial metabolism and interferes with RNA synthesis [51]. Allicin reacts with enzyme-containing thiol, thus inhibiting the Acetyl Co-A forming system and disturbing DNA synthesis and protein synthesis [52]. The present findings agree with Babu et al. [53], who reported that the MIC of *E. coli* was 1:1,000 but lower than the results obtained by Banerjee and Sarkar [54], in which the MIC was 10 mg ml⁻¹ for *E. coli*.

Cinnamon showed complete growth inhibition of all the tested *E. coli* isolates (100%) at concentrations of 125, and 63 ppm. These findings were strengthened by those described by Lu et al. [55], who investigated the

antibacterial action of cinnamon oil on *E. coli* isolates' viability and verified that their MIC ranged from 100 to 400 ppm; nonetheless, the present result was lower than that obtained by Duan and Zhao [56], in which the MIC of cinnamon against *E. coli* was 1000 ppm.

The antibacterial property of cinnamon EO may be attributed to its richness in cinnamaldehyde (the foremost constituent of cinnamon EO, comprising 85%). This aromatic aldehyde exhibits antibacterial properties against a wide variety of pathogenic microorganisms [14]. Cinnamaldehyde is a great electro-negative compound that can interrupt the biological mechanisms, including electron transport due to containing the carbonyl group that can bind to metal ions, sulfhydryl groups, and nitrogen-containing components such as proteins and nucleic acids, inhibiting the action of amino acid decarboxylases and consequently inhibiting the microorganisms' growth [57]. Cinnamon oil contains benzoic acid, benzaldehyde, and cinnamic acid, which have a lipophilic property that is responsible for its antibacterial activity as they interfere with the lipid bilayer of the cytoplasmic membrane of the bacteria, causing it to lose its integrity [58]. This backs up the idea that EOs are effective antimicrobials against a wide range of pathogenic bacteria. Finally, differences in the antibacterial findings of EO and their MICs reported in different papers could be attributable to different methodologies used to determine the MIC. Also, the purity and stability of the oil may have something to do with the different results [59].

The quantitative RT-PCR results for the *E. coli fimH* virulence gene revealed that the fold changes after treatment with 125 ppm allicin and 32 ppm cinnamon were 0.4061 and 0.1397, respectively, which indicated that the gene expression for *fimH* gene after treatment had decreased to 0.41 and 0.14 folds, respectively. In the same context, the fold changes after treatment of the second *E. coli* isolate with 125 ppm allicin and 32 ppm cinnamon were 0.4118 and 0.0693, respectively, and the fold changes after treatment of the third *E. coli* isolate with 125 ppm allicin and 32 ppm cinnamon were 0.3763 and 0.1768, respectively.

The quantitative RT-PCR results for the *E. coli int1* resistance gene showed that the fold changes after treatment with 125 ppm allicin and 32 ppm cinnamon were 0.6974 and 0.3099, respectively, which indicated that the gene expression for the *Int1* gene after treatment had decreased to 0.70 and 0.31 folds, respectively. In the same context, the fold changes after-treatment of the second *E. coli* isolate with 125 ppm allicin and 32 ppm cinnamon were 0.6417 and 0.2755, respectively, and the fold changes after treatment of the third *E. coli* isolate with 125 ppm allicin and 32 ppm cinnamon were 0.5105 and 0.2755, respectively.

The quantitative RT-PCR results for the *E. coli luxS* virulence gene showed that the fold changes after treatment

with 125 ppm allicin and 32 ppm cinnamon were 0.7120 and 0.3121, respectively, which indicated that the gene expression for the *luxS* gene after treatment had decreased to 0.71 and 0.31 folds, in that order. In the same context, the fold changes after treatment of the second *E. coli* isolate with 125 ppm allicin and 32 ppm cinnamon were 0.6156 and 0.1934, respectively, and the fold changes after treatment of the third *E. coli* isolate with 125 ppm allicin and 32 ppm cinnamon were 0.8293 and 0.3368, respectively.

Alliin and cinnamon are forms of plant-derived antimicrobial agents (PDAs) that can reduce *E. coli* virulence gene expression [60]. All of the results show that allicin and cinnamon oil could be used to make *E. coli* less dangerous [60].

Conclusion

This investigation highlights the high resistance of *E. coli* against most of the tested antimicrobials; all isolates were MDR. The presence of both virulence and resistance genes among *E. coli*. Allicin and cinnamon oils have great antimicrobial activities and could be used as alternatives to synthetic antimicrobial agents.

List of abbreviations

µg, microgram; *AadB*, aminoglycoside adenylyltransferase B; Acetyl co-A, acetyle choline A; AI-2, autoinducer-2; APEC, avian pathogenic *E. coli*; ATP, adenosine triphosphate; CLSI, Clinical and Laboratory Standards Institute; Col V, colicin V; CRA, Congo Red Agar; DNA, deoxyribonucleic acid; Eos, essential oils; EPS, extracellular polymeric substance; *FimH*, fimbrial H; h, Hour; *int1*, class 1 integrons; *Iss*, increased serum survival; *luxS*, luciferase S; MDR, Multidrug resistance; MIC, minimum inhibitory concentration; PCR, Polymerase chain reaction; PDAs, Plant-derived antimicrobial agents; ppm, part per million; *qnrS*, quinolones resistance S; q-PCR, qualitative Polymerase chain reaction; QS, Quorum sensing; RLQP, Reference laboratory for veterinary quality control on poultry production; RT-PCR, real-time Polymerase chain reaction; SYBR, synergy brands; TSI, triple sugar iron agar; YESCA/ CR, Yuba Environmental Science Charter Academy|/ congo red.

Acknowledgment

The authors are thankful to the dr. Mohammed Amin Semida and dr. Khaled mohammed shokier for their assistance.

Conflict of interest

There is no conflict of interest stated by the authors.

Authors' contribution

All the authors made an equal contribution.

References

- [1] Xu Y, Liu Y, Liu Y, Pei J, Yao S, Cheng C. Bacteriophage therapy against Enterobacteriaceae. *Virologica Sinica* 2015;30 (1):11–8; <https://doi.org/10.1007/s12250-014-3543-6>
- [2] Barnes HJ, Vaillancourt JP, Gross WB. *Colibacillosis*. In: *Diseases of poultry*, 11th edition, p 56, 2003.
- [3] Collingwood C, Kemmett K, Williams N, Wigley P. Is the concept of avian pathogenic *Escherichia coli* as a single pathotype fundamentally flawed? *Front Vet Sci* 2014; 1:5; <https://doi.org/10.3389/fvets.2014.00005>
- [4] Dziva F, Stevens MP. *Colibacillosis* in poultry. Unraveling the molecular basis of virulence of avian pathogenic *E. coli* in their natural hosts. *Avian Pathol* 2008; 37(4):355–66; <https://doi.org/10.1080/03079450802216652>
- [5] Saif YM, Barnes HJ, Glisson JR, Fadly AM, Dougland LR, Swayne DE. *Diseases of poultry*, 11th edition, Press Iowa State, Ames, IA, pp 562–6, 2003.
- [6] Phillips I, Casewell M, Cox T, de Groot B, Fiis C, Jones R, et al. Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data. *J Antimicrob Chemother* 2004; 53:28–52; <https://doi.org/10.1093/jac/dkg483>
- [7] Radwan I, Abd El-Halim M, Abed A. Molecular characterization of antimicrobial-resistant *Escherichia coli* isolated from broiler chickens. *J Vet Med Res* 2021; 27(2):128–42.
- [8] Carniello V, Peterson BW, van der Mei HC, Busscher HJ. Physicochemistry from initial bacterial adhesion to surface-programmed biofilm growth. *Adv Colloid Interface Sci* 2018; 261:1–14; <https://doi.org/10.1016/j.cis.2018.10.005>
- [9] Ostapska H, Howell PL, Sheppard DC. Deacetylated microbial biofilm exopolysaccharides: it pays to be positive. *PLoS Pathog* 2018; 14(12):1007411; <https://doi.org/10.1371/journal.ppat.1007411>
- [10] Sharma D, Misba L, Khan AU. Antibiotics versus biofilm: an emerging battleground in microbial communities. *Antimicrob Resist Infect Control* 2019; 8(76):1–10; <https://doi.org/10.1186/s13756-019-0533-3>
- [11] Shan B, Cai YZ, Brooks JD, Corke H. The *in vitro* antibacterial activity of dietary spice and medicinal herb extracts. *Int J Food Microbiol* 2007; 117(1):112–9; <https://doi.org/10.1016/j.ijfoodmicro.2007.03.003>
- [12] Radwan IA, Abed AH, Abdallah AS. Antifungal effect of carvacrol on fungal pathogens isolated from broiler chickens. *Assiut Vet Med J* 2018; 64(157):11–7; <https://doi.org/10.21608/avmj.2018.168896>
- [13] Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils: a review. *Food Chem Toxicol* 2008; 46:446–75; <https://doi.org/10.1016/j.fct.2007.09.106>
- [14] Ooi LSM, Li Y, Kam SL, Wang H, Wong EYL, Ooi VEC. Antimicrobial activities of cinnamon oil and cinnamaldehyde from the Chinese medicinal herb *Cinnamomum cassia* Blume. *Am J Chin Med* 2006; 34(03):511–22; <https://doi.org/10.1142/S0192415X06004041>
- [15] Lu X, Samuelson DR, Rasco BA, Konkel ME. Antimicrobial effect of diallyl sulphide on *Campylobacter jejuni* biofilms. *J Antimicrob Chemother*, 2012; 67(8):1915–26; <https://doi.org/10.1093/jac/dks138>
- [16] Nwachukwu ID, Slusarenko AJ. Thiosulfinates, organic polysulfanes, and related compounds: from an unusual chemistry toward a wealth of potential applications. In: *Recent advances in redox active plant and microbial products*, pp 265–88, 2014; https://doi.org/10.1007/978-94-017-8953-0_10
- [17] Quinn PJ, Markey BK, Carter ME, Donnelly WJC, Leonard FC, Maguire D. *Veterinary microbiology and microbial diseases*. 1st edition, Published Blackwell Science, Hoboken, NJ, 2002.
- [18] Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing; Twenty-Seven Informational Supplement. 2018; 37(1):M100–S27.
- [19] Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012; 18(3):268–81; <https://doi.org/10.1111/j.1469-0691.2011.03570.x>
- [20] Zhou Y, Smith DR, Hufnagel DA, Chapman MR. Experimental manipulation of the microbial functional amyloid called curli. *Methods Mol Biol* 2013; 966:53–75; https://doi.org/10.1007/978-1-62703-245-2_4
- [21] Jeff-Agboola YA, Onifade AK, Akinyele BJ, Osho IB. *In vitro* antifungal activities of essential oil from Nigerian medicinal plants against toxigenic *Aspergillus flavus*. *J Med Plants Res* 2012; 6(23):4048–56; <https://doi.org/10.5897/JMPR12.525>
- [22] Kim YB, Yoon MY, Ha JS, Seo KW, Noh EB, Son SH, Lee YJ. Molecular characterization of avian pathogenic *Escherichia coli* from broiler chickens with colibacillosis. *Poultry Sci* 2020; 99(2):1088–95; <https://doi.org/10.1016/j.psj.2019.10.047>
- [23] Awad AM, El-Shall NA, Khalil DS, El-Hack A, Mohamed E, Swelum AA, et al. Incidence, pathotyping, and antibiotic susceptibility of avian pathogenic *Escherichia coli* among diseased broiler chicks. *Pathogens* 2020; 9(2):114; <https://doi.org/10.3390/pathogens9020114>
- [24] Amer MM, Mekky HM, Amer AM, Fedawy HS. Antimicrobial resistance genes in pathogenic *Escherichia coli* isolated from diseased broiler chickens in Egypt and their relationship with the phenotypic resistance characteristics. *Vet World* 2018; 11(8):1082; <https://doi.org/10.14202/vetworld.2018.1082-1088>
- [25] El Behiry A. Diagnostic aspects and novel approach for treatment of antibiotic-resistant bacteria isolated from diarrheal calves using silver, gold and copper nanoparticles. *J Bacteriol Parasitol* 2014; 5(4):1–9; <https://doi.org/10.4172/2155-9597.1000195>
- [26] Salyers AA, Gupta A, Wang Y. Human intestinal bacteria as reservoirs for antibiotic resistance genes. *Trends Microbiol* 2004; 12(9):412–6; <https://doi.org/10.1016/j.tim.2004.07.004>
- [27] Cepas V, López Y, Muñoz E, Rolo D, Ardanuy C, Martí S, et al. Relationship between biofilm formation and antimicrobial resistance in gram-negative bacteria. *Microb Drug Resist* 2019; 25(1):72–9; <https://doi.org/10.1089/mdr.2018.0027>
- [28] Hall CW, Mah T. Molecular mechanisms of biofilm-based antibiotic resistance and tolerance in pathogenic bacteria. *FEMS Microbiol Rev* 2017; 41:276–301; <https://doi.org/10.1093/femsre/fux010>
- [29] Reichhardt C, Jacobson AN, Maher MC, Wang J, McCrate OA, Eckart M, et al. Congo red interactions with curli-producing *E. coli* and native curli amyloid fibers. *PLoS One* 2015; 10(10):e0140388; <https://doi.org/10.1371/journal.pone.0140388>
- [30] Raheel IAR, Hassan WH, Salem SSR, Salam HSH. Biofilm forming potentiality of *Escherichia coli* isolated from bovine endometritis and their antibiotic resistance profiles. *J Adv Vet Anim Res* 2020; 7(3):442–51; <https://doi.org/10.5455/javar.2020.g440>
- [31] Wang Y, Yi L, Wang Y, Wang Y, Cai Y, Zhao W, et al. Isolation, phylogenetic group, drug resistance, biofilm formation, and adherence genes of *Escherichia coli* from poultry in central China. *Poultry Sci* 2016; 95(12):2895–901; <https://doi.org/10.3382/ps/pew252>
- [32] Zając M, Sztromwasser P, Bortolaia V, Leekitcharoenphon P, Cavaco LM, Ziętek-Barszcz A, et al. Occurrence and characterization of mcr-1-positive *Escherichia coli* isolated from food-producing animals in Poland, 2011–2016. *Front Microbiol* 2019; 10:1753; <https://doi.org/10.3389/fmicb.2019.01753>
- [33] Kim T-E, Jeong Y-W, Cho S-H, Kim S-J, Kwon H-J. Chronological study of antibiotic resistances and their relevant genes in Korean avian pathogenic *Escherichia coli* isolates. *J Clin Microbiol* 2007; 45(10):3309; <https://doi.org/10.1128/JCM.01922-06>

- [34] Nhung NT, Chansiripornchai N, Carrique-Mas JJ. Antimicrobial resistance in bacterial poultry pathogens: a review. *Front Vet Sci* 2017; 4:126; <https://doi.org/10.3389/fvets.2017.00126>
- [35] Kashif J, Buriro R, Memon J, Yaqoob M, Soomro J, Dongxue D, et al. Detection of class 1 and 2 integrons, β -lactamase genes and molecular characterization of sulfonamide resistance in *Escherichia coli* isolates recovered from poultry in China. *Pak Vet J* 2013; 33(3):321–24.
- [36] Li XZ. Quinolone resistance in bacteria: emphasis on plasmid mediated mechanisms. *Int J Antimicrob Agents* 2005; 25:453–63; <https://doi.org/10.1016/j.ijantimicag.2005.04.002>
- [37] Ibrahim RA, Cryer TL, Lafi SQ, Basha E-A, Good L, Tarazi YH. Identification of *Escherichia coli* from broiler chickens in Jordan, their antimicrobial resistance, gene characterization and the associated risk factors. *BMC Vet Res* 2019; 15(1):1–16; <https://doi.org/10.1186/s12917-019-1901-1>
- [38] Tran T, Checkley S, Caffrey N, Mainali C, Gow S, Agunos A, et al. Genetic characterization of AmpC and extended-spectrum beta-lactamase (ESBL) phenotypes in *Escherichia coli* and *Salmonella* from Alberta broiler chickens. *Front Cell Infect Microbiol* 2021; 11:166; <https://doi.org/10.3389/fcimb.2021.622195>
- [39] Szmolka A, Wami H, Dobrindt U. Comparative genomics of emerging lineages and mobile resistomes of contemporary broiler strains of *Salmonella Infantis* and *E. coli*. *Front Microbiol* 2021; 12:342; <https://doi.org/10.3389/fmicb.2021.642125>
- [40] Schwartz DJ, Kalas V, Pinkner JS, Chen SL, Spaulding CN, Dodson KW, et al. Positively selected *FimH* residues enhance virulence during urinary tract infection by altering *FimH* conformation. *Proc Natl Acad Sci USA* 2013; 110(39):15530–7; <https://doi.org/10.1073/pnas.1315203110>
- [41] Wu G, Ehrlich R, Mafura M, Stokes M, Smith N, Pritchard GC, et al. *Escherichia coli* isolates from extraintestinal organs of livestock animals harbour diverse virulence genes and belong to multiple genetic lineages. *Vet Microbiol* 2012; 160(1–2):197–206; <https://doi.org/10.1016/j.vetmic.2012.05.029>
- [42] Johnson TJ, Wannemuehler, YM, Nolan LK. Evolution of the *iss* gene in *Escherichia coli*. *Appl Environ Microbiol* 2008; 74(8):2360–9; <https://doi.org/10.1128/AEM.02634-07>
- [43] Johnson TJ, Siek KE, Johnson SJ, Nolan LK. DNA sequence of a ColV plasmid and prevalence of selected plasmid-encoded virulence genes among avian *Escherichia coli* strains. *J Bacteriol* 2006; 188:745–58; <https://doi.org/10.1128/JB.188.2.745-758.2006>
- [44] Moon BM, Won GY, Choi YY, Jin JK, Oh IJ, Park JH, et al. Isolation and characteristics of avian pathogenic *Escherichia coli* from birds associated with colibacillosis. *Cairo University, Giza, Egypt*, pp 26–9, 2006.
- [45] De Kievit T, Iglewski B. Bacterial quorum sensing in pathogenic relationships. *Infect Immun* 2000; 68:4839–49; <https://doi.org/10.1128/IAI.68.9.4839-4849.2000>
- [46] Bratu S, Gupta J, Quale J. Expression of the *las* and *rhl* quorum sensing systems in clinical isolates of *Pseudomonas aeruginosa* does not correlate with efflux pump expression or antimicrobial resistance. *J Antimicrob Chemother* 2006; 58(6):1250–3; <https://doi.org/10.1093/jac/dkl407>
- [47] Davies J, Davies D. Origins and evolution of antibiotic resistance. *Microbiol Mol Biol Rev* 2010; 74(3):417–33; <https://doi.org/10.1128/MMBR.00016-10>
- [48] Wang Y, Yi L, Zhang Z, Fan H, Cheng X, Lu C. Overexpression of *luxS* cannot increase autoinducer-2 production, only affect the growth and biofilm formation in *Streptococcus suis*. *Sci World J* 2013; 924276; <https://doi.org/10.1155/2013/924276>
- [49] Niu C, Robbins CM, Pittman KJ, Osborn, JoDi L, Stubblefield BA, Simmons RB, et al. *LuxS* influences *Escherichia coli* biofilm formation through autoinducer-2-dependent and autoinducer-2-independent modalities. *FEMS Microbiol Ecol* 2013; 83(3):778–91; <https://doi.org/10.1111/1574-6941.12034>
- [50] Nazzaro F, Fratianni F, De Martino L, Coppola R, De Feo V. Effect of essential oils on pathogenic bacteria. *Pharmaceuticals* 2013; 6(12):1451–74; <https://doi.org/10.3390/ph6121451>
- [51] Ansari IA, Khan I, Khan MS, Akhtar MS. Organosulfur compounds of garlic as potent chemotherapeutic agents against cancer. In: *Anticancer plants: clinical trials and nanotechnology* pp 253–72, 2017; https://doi.org/10.1007/978-981-10-8216-0_9
- [52] Budhathoki S, Basnet A. Study of antibacterial activity of household spices. *Int J Sci Eng Res* 2018; 9(8):1700–4.
- [53] Babu AJ, RupaSundari A, Indumathi J, Srujan RVN, Sravanthi M. Study on the antimicrobial activity and minimum inhibitory concentration of essential oils of spices. *Vet World* 2011; 4(7):311; <https://doi.org/10.5455/vetworld.4.311>
- [54] Banerjee M, Sarkar PK. Inhibitory effect of garlic on bacterial pathogens from spices. *World J Microbiol Biotechnol* 2003; 19(6):565–9; <https://doi.org/10.1023/A:1025108116389>
- [55] Lu F, Ding Y, Ye X, Ding Y. Antibacterial effect of cinnamon oil combined with thyme or clove oil. *Agric Sci China* 2011; 10(9):1482–7; [https://doi.org/10.1016/S1671-2927\(11\)60142-9](https://doi.org/10.1016/S1671-2927(11)60142-9)
- [56] Duan J, Zhao Y. Antimicrobial efficiency of essential oil and freeze-thaw treatments against *Escherichia coli* O157:H7 and *Salmonella enterica* Ser. Enteritidis in strawberry juice. *Food Sci* 2009; 74(3):M131–7; <https://doi.org/10.1111/j.1750-3841.2009.01094.x>
- [57] Gupta C, Garg AP, Uniyal RC, Kumari A. Comparative analysis of the antimicrobial activity of cinnamon oil and cinnamon extract on some food-borne microbes. *Afr J Microbiol Res* 2008; 2(9):247–51; <https://doi.org/10.5897/AJMR.9000180>
- [58] Zhu H, Du M, Fox L, Zhu M-J. Bactericidal effects of Cinnamon cassia oil against bovine mastitis bacterial pathogens. *Food Control* 2016; 66: 291–9; <https://doi.org/10.1016/j.foodcont.2016.02.013>
- [59] Navarro M, Stanley R, Cusack A, Sultanbawa Y. Combinations of plant-derived compounds against *Campylobacter in vitro*. *J Appl Poult Res* 2015; 24(3):352–63; <https://doi.org/10.3382/japr/pfv035>
- [60] Baskaran SA, Kollanoor-Johny A, Nair MS, Venkitanarayanan K. Efficacy of plant-derived antimicrobials in controlling enterohemorrhagic *Escherichia coli* virulence *in vitro*. *J Food Prot* 2016; 79(11):1965–70; <https://doi.org/10.4315/0362-028X.JFP-16-104>
- [61] Frana TS, Carlson SA, Griffith RW. Relative distribution and conservation of genes encoding aminoglycoside-modifying enzymes in *Salmonella enterica* serotype Typhimurium phage type DT104. *Appl Environ Microbiol* 2001; 67:445–8; <https://doi.org/10.1128/AEM.67.1.445-448.2001>
- [62] Ghanbarpour R, Salehi M. Determination of Adhesin encoding genes in *Escherichia coli* isolates from omphalitis of chicks. *Am J Anim Vet Sci* 2010; 5(2):91–6; <https://doi.org/10.3844/ajavsp.2010.91.96>
- [63] Yaguchi K, Ogitani, T, Osawa R, Kawano M, Kokumai N, Kaneshige T, et al. Virulence factors of avian pathogenic *Escherichia coli* strains isolated from chickens with colisepticemia in Japan. *Avian Dis* 2007; 51(3):656–62; [https://doi.org/10.1637/0005-2086\(2007\)51\[656:VFOAPE\]2.0.CO;2](https://doi.org/10.1637/0005-2086(2007)51[656:VFOAPE]2.0.CO;2)
- [64] Wang X, Li S, Lu X, Hu P, Chen H, Li Z, et al. Rapid method of *luxS* and *pfs* gene inactivation in enterotoxigenic *Escherichia coli* and the effect on biofilm formation. *Mol Med Rep* 2016; 13(1):257–64; <https://doi.org/10.3892/mmr.2015.4532>
- [65] Robicsek A, Strahilevitz J, Sahn DF, Jacoby GA, Hooper DC. *qnr* prevalence in ceftazidime-resistant Enterobacteriaceae isolates from the United States. *Antimicrob Agents Chemother* 2006; 50(8):2872–4; <https://doi.org/10.1128/AAC.01647-05>
- [66] Tivendale AK, Joanne LA, Carol AG, Brendan SC, Glenn FB. Association of *iss* and *iucA*, but not *tsh*, with plasmid mediated virulence of avian pathogenic *E. coli*. *Infect Immun* 2004; 6554–60; <https://doi.org/10.1128/IAI.72.11.6554-6560.2004>